

**Determination of posterior tibia slope and slope deterioration with osteoarthritis: A radiological study in an African population.**

**Dr. Jamlick Micheni Muthuuri, MBChB, MMED (surg.), H.Dip.Orth (SA), FCS (ECSA).  
Consultant Orthopaedic Surgeon, Mombasa, Kenya.**

**Abstract**

**Study Design:** Investigational, case control study.

**Objective:** This study was performed partly to determine the mean posterior slope in the African population and partly to study slope behaviour in arthritic degeneration.

**Summary of Background:** The posterior inclination of the tibia plateaux relative to the longitudinal axis of the tibia is referred to as the posterior tibia slope (PTS). Posterior tibial slope is an important player in the kinematics of the knee joint. The depth of the slope affects the pre and post-operative range of motion (ROM). Fore knowledge of the mean PTS in a given population is important in order to keep it as close to the normal range as possible during knee replacement. The PTS is also reported to affect the integrity of both cruciate ligaments, but perhaps more the anterior cruciate ligament. Theoretically, slope deterioration should lead to anterior cruciates strain and eventual degeneration. In various studies and using diverse methods the slope has been found to range in adults from 0 to 18 degrees. Most of these studies have been done among the Caucasians, Asiatic or racial mixtures. There is paucity of information regarding PTS in the indigenous African population.

**Methods:**

Two hundred twenty four X-ray films (224) from 152 patients were used for the purpose of this study. Orthogonal radiographs for the full-limb including the knee were obtained using a standardized technique. The longitudinal axis of the tibia was obtained by manual goniometric positioning in order to draw the line marking the tibial shaft anatomical axis (TSAA). Another line was drawn tangentially over the center of the tibia plateaux joining the lowest points anteriorly and posteriorly. We measured the angle between the tangent to the tibial plateau and the perpendicular direction to TSAA. This angle was considered the posterior tibia slope. All the measurements were carried out solely by the author. The anteroposterior view was useful in grading of osteoarthritis according to the Kellgren-Lawrence classification.

## **Results:**

There were a total of 152 cases by the end of the study, 79 had non-arthritic knees (group 1) while 73 knees were osteoarthritic (group 2). Group 1 had a mean age of 48 years (95% CI 44.9 - 49.1 (2.1)) and consisted of 37 males (47%) and 42 females (53%). Group 2 had a mean age of 53 years (95% CI 45.8- 50.2 (2.2)) and consisted of 33 males (47%) and 40 females (53%). Control group had a mean slope angle of 6.8 compared to the arthritic group mean of 6.1 (95% confidence level  $P < 0.0505$ ) which was significant.

## **Conclusion:**

Our findings are that the mean posterior tibia slope among the African population is about  $7^\circ$ . We also conclude that the slope decreases with osteoarthritic degeneration. This would also suggest cruciate insufficiency that would result in either anterior or posterior tibia instability.

## **Introduction**

The posterior inclination of the tibia plateaux relative to the longitudinal axis of the tibia is referred to as the posterior tibia slope (PTS). The posterior tibial slope plays a very important role in the kinematics and biomechanics of the knee joint. It has been shown that increasing PTS increases the tibial shear force and anterior tibial translation at the knee [1]. This is particularly important in individuals undergoing proximal tibial osteotomy as these forces will accelerate degenerative changes. This altered PTS will affect not only the integrity of the cruciate ligaments but also the range of motion (ROM). Increased medial PTS has been reported to be associated with increased anterior cruciate ligament (ACL) injuries due to increased anterior tibia translation [2, 3]. Although some authors have reported increased stability of posterior cruciate ligament deficient knees with deepened posterior tibial slopes [4]. Therefore, the slope should be maintained or corrected to maintain normal kinematics. These knee forces are highly significant in osteoarthritis and in the survival and function of knee arthroplasty; with potential complications, such as progressive loosening of implants, fractures of polyethylene inserts and wears [5, 6].

In degenerative joint disease, the slope and other aspects of the tibial plateaux are modified by bone and cartilage wear. The PTS will then measure the sum of these changes and may not reflect accurately the normal slope and mechanical axis of the knee [7]. The slope is reported to deepen in

osteoarthritis; meaning increased articular surface contact and increased tibial translation [8]. Total knee replacement aims to restore the mechanical axis of the natural knee joint. This axis will be changed by an altered PTS; yet after replacement the PTS is expected to play a key role in crucial movements, particularly in achieving optimum knee flexion [9, 10].

In various studies using diverse methods, the slope has been found to range in adults from 0 to 18 degrees [11, 12]. Most of these studies are among the Caucasians, Asiatic or racial mixtures. The PTS in the African population has not been determined or documented.

### **MATERIAL AND METHODS:**

We conducted measurements of posterior tibia slopes on knees in patients of African origin aged 40 years and above. These were patients requiring knee radiographs for various reasons, chiefly knee pain. Patients willingly consented for the use of their radiographs for the study. Two hundred and twenty four films (224) were examined from one hundred and fifty two patients (152). Radiographs of both knees were used when available and comparable; the mean value between the two was recorded. Orthogonal radiographs for the full-limb including the knee were obtained using a standardized technique.

A true lateral radiograph in extension with the exact superimposition of the femoral condyles was done in order to obtain various morphometric measurements. The radiograph was considered adequate if it exposed the knee and at least two thirds of the proximal tibia. The longitudinal axis of the tibia was obtained by manual goniometric positioning in order to draw the line marking the tibial shaft anatomical axis (TSAA). Another line was drawn tangentially over the center of the tibia plateaux joining the lowest points anteriorly and posteriorly. We measured the angle between the tangent to the tibial plateau and the perpendicular direction to TSAA. This angle was considered the posterior tibia slope (Fig.1). All the measurements were carried out solely by the author.



Fig. 1 shows how the PTS is determined. Line 1 is the perpendicular direction to TSAA while line 2 is drawn tangentially over the center of the tibia plateaux.

The X-ray films done in the anteroposterior view was useful in grading of osteoarthritis. The patients were grouped into two; those with non-arthritic knees and those with overt osteoarthritis (Kellgren-Lawrence >2). The following patients were excluded from the study. All those found to have fractures, severe osteoporosis, tumours, Charcot's disease or osteolysis from any cause.

### Statistical Analysis

The results were analysed using the IBM SPSS statistic version 19 and computer generated statistic formulas. Statistical analysis included both descriptive and inferential measures. Mean values and standard deviations were calculated for standard valuables, and frequent counts and percentages were obtained for discrete valuables. All confidence intervals (designated CI) are at 95% confidence level.  $X^2$  test was used for cross tabulation analysis. For continuous valuable (e.g. patient age), the 2 sample independent T test was used.

The sample size was calculated at 90% power in order to demonstrate any effect between the slope in arthritic and non-arthritic knees.

### RESULTS:

The posterior tibial slope was obtained from 224 films from 152 patients. There were 70 males out of whom 33 (47%) had arthritic knees. There were 82 females out of whom 40 (49%) had arthritic knees, Table 1.


Table 1 Sample distribution

Group	Male	Female	Total
Arthritic (Group 1)	33	40	73

Normal (Group 2)	37	42	79
Total	70	82	152

There was equal distribution of cases between age groups. Patients' age did not significantly differ between genders in both groups ( $P > 0.181$ ). However, 71% of the patients were aged between 40 and 60 years (Table.2 and Fig. 2).


Table.2 and Fig. 2: Age distribution



As expected there is exponential increase in osteoarthritis with age. Table 3 and Fig.3 confirm this expectation.

Table 3 and Fig.3: Age groups and Arthritis.

AGE	Number	Normal	Arthritic	% Arthritic
40-49	55	38	17	31
50-59	53	25	28	53
60-69	23	11	12	52
>70	21	5	16	76
<b>TOTAL</b>	<b>152</b>	<b>79</b>	<b>73</b>	<b>48</b>



The range for posterior tibia slope was found to be 3° to 15° (Table 4 & Fig. 4). The graph is skewed to the left. The mean slope angle was found to be 6.6° (95% confidence limits of 6.2° to 7.0° (0.4 confidence interval)). The arthritic group had a mean of 6.1° against control group mean of 6.8°. This was found to be significant (P>0.0505). This showed deterioration of the slope by 0.7° (Table 5).

Table 4 Slope measurements

SLOPE	NORMAL KNEES	ARTHRITIC KNEES	TOTAL
3°	3	4	7
4°	7	8	15
5°	19	16	35
6°	14	21	35
7°	14	12	26
8°	6	1	7
9°	3	5	8

10°	3	2	5
11°	2	2	4
12°	3	1	4
13°	0	0	0
14°	5	2	7
<b>MEAN</b>	<b>79</b>	<b>73</b>	<b>152</b>

Fig. 4 Slope measurements




Table.5. Slope changes with Arthritis

<b>MEANS</b>	<b>NORMAL</b>	<b>ARTHRITIC</b>	<b>SAMPLE</b>	<b>CI</b>	<b>S.e.d</b>	<b>SD</b>	<b>Z SCORE</b>	<b>P VALUE</b>
	<i>m0</i>	<i>m1</i>	<i>m</i>					
<b>MALE</b>	6.7	6	6.6	0.67	0.81342	2.8	<b>-0.861</b>	<b>0.1949</b>
<b>FEMALE</b>	6.4	6.1	6.2	0.40	0.398538	1.8	<b>-0.753</b>	<b>0.2266</b>
<b>SAMPLE</b>	6.8	6.1	6.6	0.40	0.412151	2.5	<b>-1.643</b>	<b>0.0505</b>

## **DISCUSSION:**

The normal values for the posterior tibial plateau slope have not been fully mapped out for all populations. However, what is reported show a high variability between races and ranges from 0° to 18°.

The posterior tibial slope has been assessed using various methods, ranging from direct cadaveric measurements to CT scans and MRI. Various methods of assessment on plain radiographs have been described. Brazier et al described the different anatomical axes against which the PTS could be measured [13]. The lines included the tibial proximal anatomical axis (TPAA); tibial shaft anatomical axis (TSAA); posterior tibial cortex (PTC); anterior tibial cortex (ATC); fibular proximal anatomical axis (FPAA) and fibular shaft axis (FSA). They concluded that among the proximal axes, the TPAA and PTC gave higher reliability.

The ideal way to measure the PTS is against the anatomical axis of the tibia (TSAA). This will require a radiograph of the whole tibia which is usually not taken routinely in clinical practice. The tibial proximal anatomical axis (TPAA) does not exactly match the mechanical axis of the tibia but gives a close correlation. We used the TSAA in this study.

Computer Tomographic scan and MRI scan images have also been used for measuring the slope on medial and lateral sides separately [11, 14]. In a MRI study with separate measurement for medial and lateral plateaux slopes, the mean posterior slope was higher in the lateral plateau than medial [11]. In this study, we found the slope varied from 3°-14° averaging 6.6°. There was a higher slope in males than in females (mean of 6.6 vs 6.2). This is in contrast to the findings of Hashemi who found slopes in females higher than males [11]. Yoga R et al found the mean preoperative posterior tibial plateau slope in Malaysian population to be 10.0 degrees, 9.9° in Chinese and 8.8° in Indians [8], while Chiu et al, noted that the average posterior plateau slope in Chinese patients was 11.5 degrees [15]. Haddad et al, found mean PTS of 4.2° in Whites, 7.9° in Asians and 6.0° in Afro-Caribbean/European [16]. This group found these racial differences to be statistically significant ( $p < 0.001$ ).

Didia BC et al performed a radiological study on 212 lateral tibial radiographs measuring PTS with reference to the ATC found a mean PTS of 12.3° [17].



Some authors have reported a greater incidence of ACL rupture when the PTS angle is increased [18]. Female athletes have been found to have a higher incidence of ACL injuries (up to nine times more common)[19, 20] and also to be more susceptible to non-contact ACL injuries. Theoretically, therefore, increased PTS in certain populations might be associated with increased stress on the ACL. This increased strain or incompetence of the ACL will allow the tibia to sublux anteriorly. The effect is a reversal of the posterior slope which will then tend to flatten. A careful look at the knees during replacement surgery shows deepening concavities of the central part of the tibial condyle leaving the rim as protected by the meniscus almost intact. However, as stated earlier, the slope has been reported to deepen in osteoarthritis; meaning increased articular surface contact and increased tibial translation [8]. Chiu et al also concluded that osteoarthritis increase the slope by two to three degrees [15]. Our study show that the slope flattens as osteoarthritis progresses (6.8 vs 6.1) with a p value=0.0505. Our patients were in all grades of arthritis from moderate to severe (Kellgren & Lawrence grade II-IV). Further studies with large samples dealing with end stage osteoarthritis (Kellgren & Lawrence grade IV) will perhaps, settle this matter.

## **CONCLUSIONS:**

Our findings are that the mean posterior tibia slope among the African population is 6°. We also conclude that the slope decreases with osteoarthritic degeneration. This would also suggest cruciate degeneration and insufficiency that would result in either anterior or posterior tibia instability.

## **CLINICAL RELEVANCE:**

These results suggest that the PTS in the African is around 7°. This information is important during arthroplasty. Secondly, degenerative osteoarthritis affects tibia slope. A larger study is required to establish whether the slope increases or decreases. Lastly, it is important to maintain PTS during high tibial osteotomy; as again increase or decrease will affect long term stability.

## **REFERENCES**

1. Shelburne KB, K.H., Sterett WI, Pandya MG., *Effect of posterior tibial slope on knee biomechanics during functional activity.* . Orthop Res., 2011. **29**(2): p. 223-31.
2. Brandon ML, H.P., Bonamo JR et al. , *The association between posterior-inferior tibial slope and anterior cruciate ligament insufficiency.* . Arthroscopy 2006 ; , 2006(22 ): p. 894-849.
3. Todd MS, L.S., Garcia E, DeBerardino TM, Cameron KL. , *The relationship between posterior tibial slope and anterior cruciate ligament injuries.* . Am J Sports Med 2010. **38**(2010): p. 63-67.
4. Giffin JR, S.K., Zantop T, Vogrin TM, Woo SL, Harner CD., *Importance of tibial slope for stability of the posterior cruciate ligament deficient knee.* . Am J Sports Med. , 2007. **35**(9): p. 1443-9.
5. Stijak L, B.Z., Kadija M, Stanković G, Djulejić V, Milovanović D, Filipović B. , *The role of the posterior tibial slope on rupture of the anterior cruciate ligament.* . J Am Acad Orthop Surg., 2013. **21**(1): p. 41-50.
6. Migaud H, D.L.A., Dohin B, Cloutier JM, Gougeon F, Duquennoy A. , *Influence of the tibial slope on tibial translation and mobility of non-constrained total knee prosthesis.* . Rev Chir Orthop Reparatrice Appar Mot. , 1996. **82**(1): p. 7-13.
7. Jenny JY, B.C., Ballonzoli L, Meyer N., *Difficulties and reproducibility of radiological measurement of the proximal tibial axis according to Lévine].* Knee Surg Sports Traumatol Arthrosc. , 2008. **16**(4): p. 373-7.
8. Yoga R, S.N., Suresh C., *Posterior slope of the tibia plateau in malaysian patients undergoing toatal knee replacement.* Malaysian Orthopaedic Journal, 2009. **3**(1): p. 78-80.
9. Malviya A, L.E., Weir DJ, Deehan DJ. , *Predicting range of movement after knee replacement : the importance of posterior condylar offset and tibial slope.* . Knee Sur Sports Traumatol Arthrosc 2009(17): p. 491-498.

10. Bellemans J, R.F., Duerinckx J, Banks S, Vandenuecker H. , *The influence of tibial slope on maximal flexion after total knee arthroplasty.* . *Traumatol Arthrosc* 2005 ; , 2005(13 : ): p. 193-196.
11. Hashemi J, C.N., Gill B et al. , *The geometry of the tibial plateau and its influence on the biomechanics of the tibiofemoral joint.* . *J Bone Joint Surg* 2008 ; , 2008(90-A : ): p. 2724-2734.
12. Genin P, W.G., Julliard R, *The tibial slope. Proposal for a measurement method.* . *Journal of Radiology*, 1993. **74**(1): p. 27-33.
13. Brazier J, M.H., Gougeon F, Cotten A, Fontaine C, Duquennoy A, *Evaluation of methods for radiographic measurement of the tibia slope; a study of 83 healthy knees.* *Rev Chir Orthop Reparatrice Appar Mot.*, 1996. **82**(3): p. 195-200.
14. Hudek R, S.S., Regenfelder F, Fuchs B, Koch PP., *Novel measurement technique of the tibial slope on conventional MRI.* . *Clin Orthop Relat Res* 2009 ; , 2009(467 : ): p. 2066-2072.
15. Chiur Ky, Z.S., Zhang GH., *Posterior slope of tibial plateau in Chinese.* *J Arthroplasty.*, 2000. **15**(2): p. 224-227.
16. Haddad B, K.S., Mannan K, Scott G. , *Evaluation of the posterior tibial slope on MR images in different population groups using the tibial proximal anatomical axis.* . *Vojnosanit Pregl.*, 2012. **69**(10): p. 864-8.
17. Didia BC, J.B., *Posterior slope of tibial plateau in adult nigerian subjects.* . *Int J Morphology* 2009; , 2009. **27**: p. 201-204.
18. Sonnery-Cottet B, A.P., Cucurulo T et al. , *The influence of the tibial slope and the size of the inter -condylar notch on rupture of the anterior cruciate ligament.*. *J Bone Joint Surg* 2011 ; 93-B : . 2011(93-B): p. 1475-1478.

19. Cimino F, V.B., Setter D. , *Anterior cruciate ligament injury : diagnosis, management, and prevention.* . Am Fam Physician 2010 ; , 2010(82 :): p. 917-922.
20. Vauhnik R, M.M., Rutherford OM et al. , *Rate and risk of anterior cruciate ligament injury among sports -women in Slovenia.* . J Athl Train 2011. **2011**(46): p. 92-98.